





Institut für Angewandte Physik Physikalisches Institut RHEINISCHE FRIEDRICH-WILHELMS-UNI-VERSITÄT BONN

## COLLOQUIUM "OPTICS AND CONDENSED MATTER"

## Sven Höfling

University of Würzburg, Germany

## Polariton lattices in patterned microcavities

## Simon Betzold<sup>1</sup>, Christian Schneider<sup>2</sup>, Sebastian Klembt<sup>1</sup>, Sven Höfling<sup>1</sup>

 <sup>1</sup> Julius-Maximilians-Universität Würzburg & Würzburg-Dresden Cluster of Excellence ct.qmat, Physikalisches Institut, Lehrstuhl für Technische Physik, Am Hubland, 97074 Würzburg, Germany
<sup>2</sup> Institute of Physics, Carl von Ossietzky University, 26129 Oldenburg, Germany

Polaritons are quasiparticles composed partly of light and partly of matter, combine pronounced nonlinearities arising from their excitonic component with the possibility of on-chip implementation. In this presentation, techniques to confine polaritons and build up lattices will be presented, both for III-V and fluorescent protein semiconductors. We report the formation of confined mode and band structures as well as the lasing properties of exciton-polaritons in single or coupled microcavities [1-5], for instance arranged in a linear periodic 1D, Su-Schrieffer-Heeger chains, honeycomb or other lattices.

[1] S. Betzold et al., ACS Photonics 7, 384 (2020)

- [2] M. Dusel et al., Nat. Commun., 11, 2863 (2020)
- [3] M. Dusel et al., Nano Lett., 21, 6398 (2021)
- [4] T. H. Harder et al., ACS Photonics 8, 1377 (2021)
- [5] S. Klembt et al., Exciton-polariton topological insulator. Nature 562, 552 (2018)

**February 6th, starting with discussion at 17:00 h, talk at 17:15 h, live IAP lecture hall or via Zoom** <u>https://uni-bonn.zoom.us/j/98441612025?pwd=a01SSjlkY1Q3SDFhL09JQk1qc1V6dz09</u> Meeting-ID: 984 4161 2025 Kenncode: 294164